

PRX I Polyclonal Antibody

Catalog # AP63494

Product Information

Application	WB, IHC-P
Primary Accession	Q06830
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22110

Additional Information

Gene ID	5052
Other Names	PRDX1; PAGA; PAGB; TDPX2; Peroxiredoxin-1; Natural killer cell-enhancing factor A; NKEF-A; Proliferation-associated gene protein; PAG; Thioredoxin peroxidase 2; Thioredoxin-dependent peroxide reductase 2
Dilution	WB~~WB: 1:1000-2000 IHC: 1:200-500 IHC-P~~N/A
Format	PBS, pH 7.4, containing 0.09% (W/V) sodium azide as Preservative and 50% Glycerol.
Storage Conditions	-20°C


Protein Information

Name	PRDX1
Synonyms	PAGA, PAGB, TDPX2
Function	Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2) (PubMed: 9497357). Reduces an intramolecular disulfide bond in GPD5 that gates the ability to GPD5 to drive postmitotic motor neuron differentiation (By similarity).
Cellular Location	Cytoplasm. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV


Background

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H₂O₂ (PubMed:[9497357](#)). Reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation (By similarity).

Images



Western blot analysis of 1) Hela, 2) Mouse Brain, 3) Rat Brain using PRX I Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded Human Hepatocarcinoma using PRX I Polyclonal Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.