

Transferrin Monoclonal Antibody(5A4)

Catalog # AP63654

Product Information

Application	WB, IHC-P
Primary Accession	P02787
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	77064

Additional Information

Gene ID	7018
Other Names	TF; Serotransferrin; Transferrin; Beta-1 metal-binding globulin; Siderophilin
Dilution	WB~~WB 1:2000-5000, IHC 1:100-200 IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	TF (HGNC:11740)
Function	Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in stimulating cell proliferation. (Microbial infection) Serves as an iron source for parasite T.brucei (strain 427), which capture TF via its own transferrin receptor ESAG6:ESAG7 and extract its iron for its own use.
Cellular Location	Secreted.
Tissue Location	Expressed by the liver and secreted in plasma.

Background

Transferrins are iron binding transport proteins which can bind two Fe(3+) ions in association with the binding of an anion, usually bicarbonate. It is responsible for the transport of iron from sites of absorption and heme degradation to those of storage and utilization. Serum transferrin may also have a further role in

stimulating cell proliferation.

Images



Western blot analysis of Human Serum with Transferrin Mouse mAb diluted at 1:2,000.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.