

EAAT2 Polyclonal Antibody

Catalog # AP63677

Product Information

Application	WB
Primary Accession	<u>P43004</u>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62104

Additional Information

Gene ID	6506
Other Names	Excitatory amino acid transporter 2 (Glutamate/aspartate transporter II) (Sodium-dependent glutamate/aspartate transporter 2) (Solute carrier family 1 member 2)
Dilution	WB~~WB 1:1000-2000
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SLC1A2 (<u>HGNC:10940</u>)
Function	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also L-aspartate and D-aspartate (PubMed: <u>14506254</u> , PubMed: <u>15265858</u> , PubMed: <u>26690923</u> , PubMed: <u>7521911</u>). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed: <u>14506254</u>). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed: <u>14506254</u>). Essential for the rapid removal of released glutamate from the synaptic cleft, and for terminating the postsynaptic action of glutamate (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein

Background

Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L-glutamate and also

L-aspartate and D-aspartate (PubMed:<u>7521911</u>, PubMed:<u>14506254</u>, PubMed:<u>15265858</u>, PubMed:<u>26690923</u>). Functions as a symporter that transports one amino acid molecule together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion (PubMed:<u>14506254</u>). Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (PubMed:<u>14506254</u>). Essential for the rapid removal of released glutamate from the synaptic cleft, and for terminating the postsynaptic action of glutamate (By similarity).

Images



Western blot analysis of 1) Mouse Brain Tissue, 2)Rat Brain Tissue with EAAT2 Rabbit pAb diluted at 1:2,000.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.