

GABA A Receptor α 3 Polyclonal Antibody

Catalog # AP63679

Product Information

Application	WB, IHC-P
Primary Accession	<u>P34903</u>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55165

Additional Information

Gene ID	2556
Other Names	Gamma-aminobutyric acid receptor subunit alpha-3 (GABA(A) receptor subunit alpha-3)
Dilution	WB~~WB 1:1000-2000, IHC 1:100-200 IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	GABRA3 (<u>HGNC:4077</u>)
Function	Alpha subunit of the heteropentameric ligand-gated chloride channel gated by gamma-aminobutyric acid (GABA), a major inhibitory neurotransmitter in the brain (PubMed: <u>16412217</u> , PubMed: <u>29053855</u>). GABA- gated chloride channels, also named GABA(A) receptors (GABAAR), consist of five subunits arranged around a central pore and contain GABA active binding site(s) located at the alpha and beta subunit interface(s) (By similarity). When activated by GABA, GABAARs selectively allow the flow of chloride anions across the cell membrane down their electrochemical gradient (PubMed: <u>16412217</u> , PubMed: <u>29053855</u>). Chloride influx into the postsynaptic neuron following GABAAR opening decreases the neuron ability to generate a new action potential, thereby reducing nerve transmission (PubMed: <u>16412217</u> , PubMed: <u>29053855</u>).
Cellular Location	Postsynaptic cell membrane {ECO:0000250 UniProtKB:P14867}; Multi-pass membrane protein. Cell membrane {ECO:0000250 UniProtKB:P14867}; Multi-pass membrane protein

Background

GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel.

Images



Western blot analysis of 1) Mouse Brain Tissue, 2)Rat Brain Tissue with GABA A Receptor α 3 Rabbit pAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded Rat BrainTissue using GABA A Receptor α 3 Rabbit pAb diluted at 1:200.

Immunohistochemical analysis of paraffin-embedded Mouse BrainTissue using GABA A Receptor α 3 Rabbit pAb diluted at 1:200.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.