

# Cystatin C mouse Monoclonal Antibody(3B12)

Catalog # AP63751

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P01034</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Calculated MW</b>	15799

## Additional Information

---

<b>Gene ID</b>	1471
<b>Other Names</b>	CST3
<b>Dilution</b>	WB~~WB 1:1000-2000
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	CST3
<b>Function</b>	As an inhibitor of cysteine proteinases, this protein is thought to serve an important physiological role as a local regulator of this enzyme activity.
<b>Cellular Location</b>	Secreted.
<b>Tissue Location</b>	Expressed in submandibular and sublingual saliva but not in parotid saliva (at protein level). Expressed in various body fluids, such as the cerebrospinal fluid and plasma. Expressed in highest levels in the epididymis, vas deferens, brain, thymus, and ovary and the lowest in the submandibular gland


## Background

---

As an inhibitor of cysteine proteinases, this protein is thought to serve an important physiological role as a local regulator of this enzyme activity.

## Images

---



Western Blot analysis of Cystatin C protein using antibody diluted at 1:1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.