

CD10 Rabbit Polyclonal Antibody

Catalog # AP63791

Product Information

Application	IHC-P
Primary Accession	P08473
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85514

Additional Information

Gene ID	4311
Other Names	MME
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	MME {ECO:0000303 PubMed:27588448, ECO:0000312 HGNC:HGNC:7154}
Function	Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed: 15283675 , PubMed: 6208535 , PubMed: 6349683 , PubMed: 8168535). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed: 17101991 , PubMed: 6349683). Catalyzes cleavage of bradykinin, substance P and neuropeptides (PubMed: 6208535). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed: 15283675 , PubMed: 6349683). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed: 16254193 , PubMed: 2531377 , PubMed: 2972276). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed: 20876573).
Cellular Location	Cell membrane; Single-pass type II membrane protein

Background

Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids

(PubMed:[15283675](#), PubMed:[8168535](#)). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed:[17101991](#)). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed:[15283675](#)). Involved in the degradation of atrial natriuretic factor (ANF) (PubMed:[2531377](#), PubMed:[2972276](#)). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed:[20876573](#)).

Images



Immunohistochemical analysis of paraffin-embedded Human Kidney Tissue using CD10 Rabbit pAb diluted at 1:500.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.