

# CLNS1A Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6520c

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P54105</a>
<b>Other Accession</b>	<a href="#">Q04753</a> , <a href="#">Q28678</a> , <a href="#">Q61189</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rabbit, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB19566
<b>Calculated MW</b>	26215
<b>Antigen Region</b>	148-176

## Additional Information

---

<b>Gene ID</b>	1207
<b>Other Names</b>	Methylosome subunit pICln, Chloride channel, nucleotide sensitive 1A, Chloride conductance regulatory protein ICln, I(Cln), Chloride ion current inducer protein, ClCI, Reticulocyte pICln, CLNS1A, CLCI, ICLN
<b>Target/Specificity</b>	This CLNS1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 148-176 amino acids from the Central region of human CLNS1A.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CLNS1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CLNS1A
-------------	--------

## Synonyms

CLCI, ICLN

## Function

Involved in both the assembly of spliceosomal snRNPs and the methylation of Sm proteins (PubMed:[10330151](#), PubMed:[11713266](#), PubMed:[18984161](#), PubMed:[21081503](#)). Chaperone that regulates the assembly of spliceosomal U1, U2, U4 and U5 small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre- mRNAs (PubMed:[10330151](#), PubMed:[18984161](#)). Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP (Sm core) (PubMed:[10330151](#)). In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG are trapped in an inactive 6S pICln-Sm complex by the chaperone CLNS1A that controls the assembly of the core snRNP (PubMed:[10330151](#), PubMed:[18984161](#)). Dissociation by the SMN complex of CLNS1A from the trapped Sm proteins and their transfer to an SMN-Sm complex triggers the assembly of core snRNPs and their transport to the nucleus (PubMed:[10330151](#), PubMed:[18984161](#)).

## Cellular Location

Cytoplasm, cytosol. Nucleus. Cytoplasm, cytoskeleton. Note=A small fraction is also associated with the cytoskeleton (PubMed:[18984161](#))


## Background

CLNS1A is a protein that functions in multiple regulatory pathways. The protein complexes with numerous cytosolic proteins and performs diverse functions including regulation of small nuclear ribonucleoprotein biosynthesis, platelet activation and cytoskeletal organization. The protein is also found associated with the plasma membrane where it functions as a chloride current regulator.

## References



Yeung,C.H., Biol. Reprod. 73 (5), 1057-1063 (2005)

## Images



Western blot analysis of CLNS1A Antibody (Center) (Cat.# AP6520c) in HL-60 and Hela cell line lysates (35ug/lane). CLNS1A (arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human colon carcinoma reacted with CLNS1A Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of hela cells using CLNS1A Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.