

ELOVL6 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6524a

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q9H5J4
Other Accession	Q920L6 , Q920L5 , Q5ZJR8
Reactivity	Human, Mouse
Predicted	Chicken, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB19614
Calculated MW	31376
Antigen Region	32-58

Additional Information

Gene ID	79071
Other Names	Elongation of very long chain fatty acids protein 6, 3-keto acyl-CoA synthase ELOVL6, ELOVL fatty acid elongase 6, ELOVL FA elongase 6, Fatty acid elongase 2, hELO2, Fatty acyl-CoA elongase, Long-chain fatty-acyl elongase, Very-long-chain 3-oxoacyl-CoA synthase 6, ELOVL6, FACE, LCE
Target/Specificity	This ELOVL6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 32-58 amino acids from the N-terminal region of human ELOVL6.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ELOVL6 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ELOVL6 {ECO:0000255 HAMAP-Rule:MF_03206}
------	--

Synonyms	FACE, LCE
Function	Catalyzes the first and rate-limiting reaction of the four reactions that constitute the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process allows the addition of 2 carbons to the chain of long- and very long-chain fatty acids (VLCFAs) per cycle. Condensing enzyme that elongates fatty acids with 12, 14 and 16 carbons with higher activity toward C16:0 acyl-CoAs. Catalyzes the synthesis of unsaturated C16 long chain fatty acids and, to a lesser extent, C18:0 and those with low desaturation degree. May participate in the production of saturated and monounsaturated VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators.
Cellular Location	Endoplasmic reticulum membrane {ECO:0000255 HAMAP-Rule:MF_03206, ECO:0000269 PubMed:20937905}; Multi- pass membrane protein {ECO:0000255 HAMAP-Rule:MF_03206}
Tissue Location	Ubiquitous..


Background

Fatty acid elongases (EC 6.2.1.3), such as ELOVL6, use malonyl-CoA as a 2-carbon donor in the first and rate-limiting step of fatty acid elongation.


References

Lu,Y., J. Lipid Res. 49 (12), 2582-2589 (2008)


Images




Anti-ELOVL6 Antibody (N-term) at 1:1000 dilution + human lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of ELOVL6 antibody (N-term) (Cat.# AP6524a) in mouse liver tissue lysates (35ug/lane). ELOVL6 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma reacted with ELOVL6 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



ELOVL6 Antibody (N-term) (Cat.#AP6524a) flow cytometry analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.