

# FABP3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6528a

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P05413</a>
<b>Other Accession</b>	<a href="#">P07483</a> , <a href="#">O02772</a> , <a href="#">P11404</a> , <a href="#">P10790</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Bovine, Pig, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB19633
<b>Calculated MW</b>	14858
<b>Antigen Region</b>	1-30

## Additional Information

---

<b>Gene ID</b>	2170
<b>Other Names</b>	Fatty acid-binding protein, heart, Fatty acid-binding protein 3, Heart-type fatty acid-binding protein, H-FABP, Mammary-derived growth inhibitor, MDGI, Muscle fatty acid-binding protein, M-FABP, FABP3, FABP11, MDGI
<b>Target/Specificity</b>	This FABP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human FABP3.
<b>Dilution</b>	WB~~1:2000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	FABP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FABP3
-------------	-------

<b>Synonyms</b>	FABP11, MDGI
<b>Function</b>	FABPs are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters.
<b>Cellular Location</b>	Cytoplasm.


## Background

The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells.


## References

Iwayama,Y., Am. J. Med. Genet. B Neuropsychiatr. Genet. (2009)  
Lazary,A., Eur. J. Endocrinol. 159 (2), 187-196 (2008)

## Images





Western blot analysis of lysate from human heart tissue lysate, using FABP3 Antibody (N-term)(Cat. #AP6528a). AP6528a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.




Western blot analysis of FABP3 antibody (N-term) (Cat.# AP6528a) in mouse heart tissue lysates (35ug/lane). FABP3 (arrow) was detected using the purified Pab.

Formalin-fixed and paraffin-embedded human hepatocarcinoma with FABP3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of MDA-231 cells using FABP3 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Western blot analysis of FABP3 (arrow) using rabbit polyclonal FABP3 Antibody (N-term) (Cat. #AP6528a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the FABP3 gene.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.