10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# TYSY Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP6682b

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, FC, IF, E |
| **Primary Accession** | P04818 |
| **Reactivity** | Human |
| **Predicted** | Mouse, Rat, Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Calculated MW** | 35716 |
| **Antigen Region** | 265-294 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 7298 |
| **Other Names** | Thymidylate synthase, TS, TSase, TYMS, TS |
| **Target/Specificity** | This TYSY antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 265-294 amino acids from the C-terminal region of human TYSY. |
| **Dilution** | WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | TYSY Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | TYMS ( HGNC:12441) |
| **Synonyms** | TS |
| **Function** | Catalyzes the reductive methylation of 2'-deoxyuridine 5'- monophosphate (dUMP) to thymidine 5'-monophosphate (dTMP), using the cosubstrate, 5,10- |

| | |
|---|---|
| | methylenetetrahydrofolate (CH2H4folate) as a 1- carbon donor and reductant and contributes to the mitochondrial and nuclear de novo thymidylate biosynthesis pathway. |
| **Cellular Location** | Nucleus. Cytoplasm Mitochondrion. Mitochondrion matrix. Mitochondrion inner membrane. Note=Localized to the nucleus during S and G2/M phases of the cell cycle. As a component of the de novo thymidylate synthesis complex, localizes specifically to replication forks during DNA synthesis (PubMed:22235121) |


## Background

Thymidylate synthase catalyzes the methylation of deoxyuridylate to deoxythymidylate using 5,10-methylenetetrahydrofolate (methylene-THF) as a cofactor. This function maintains the dTMP (thymidine-5-prime monophosphate) pool critical for DNA replication and repair. The enzyme has been of interest as a target for cancer chemotherapeutic agents. It is considered to be the primary site of action for 5-fluorouracil, 5-fluoro-2-prime-deoxyuridine, and some folate analogs.


## References

Ren,D.N., J Surg Oncol (2009) Schiffer,C.A., Biochemistry 34 (50), 16279-16287 (1995)

## Images






All lanes: Anti-TYSY Antibody (C-term) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 33 KDa Blocking/Dilution buffer: 5% NFDM/TBST.




Formalin-fixed and paraffin-embedded human colon carcinoma with TYSY Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Flow cytometric analysis of WiDr cells using TYSY Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Confocal immunofluorescent analysis of TYSY Antibody (C-term)(Cat#AP6682b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green).Actin filaments have been labeled with Alexa Fluor? 555 phalloidin (red).

## Citations

- DNA methylation-regulated miR-193a-3p dictates resistance of hepatocellular carcinoma to 5-fluorouracil via repression of SRSF2 expression.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.