

Bcl-2 (phospho Ser70) Polyclonal Antibody

Catalog # AP66963

Product Information

Application	WB, IHC-P, IP
Primary Accession	P10415
Reactivity	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26266

Additional Information

Gene ID	596
Other Names	BCL2; Apoptosis regulator Bcl-2
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A IP~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	BCL2
Function	Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed: 1508712 , PubMed: 8183370). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed: 11368354). Appears to function in a feedback loop system with caspases (PubMed: 11368354). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed: 11368354). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during non-starvation conditions and inhibits their autophagy function (PubMed: 18570871 , PubMed: 20889974 , PubMed: 21358617). May attenuate inflammation by impairing NLRP1- inflammasome activation, hence CASP1 activation and IL1B release (PubMed: 17418785).
Cellular Location	Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250 UniProtKB:P10417}

Tissue Location

Expressed in a variety of tissues.

Background

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. Regulates cell death by controlling the mitochondrial membrane permeability. Appears to function in a feedback loop system with caspases. Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1). May attenuate inflammation by impairing NLRP1-inflammasome activation, hence CASP1 activation and IL1B release (PubMed:[17418785](#)).

Images

Western Blot analysis of various cells using Phospho-Bcl-2 (S70) Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HeLa cells using Phospho-Bcl-2 (S70) Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.