

c-Myc (phospho Thr58) Polyclonal Antibody

Catalog # AP67001

Product Information

Application WB, IHC-P, IP **Primary Accession** P01106

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW50565

Additional Information

Gene ID 4609

Other Names MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix

protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunoprecipitation: 2-5 ug/mg lysate. ELISA: 1/10000. Not yet tested in

other applications. IHC-P~~N/A IP~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name MYC

Synonyms BHLHE39

Function Transcription factor that binds DNA in a non-specific manner, yet also

specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed: 24940000, PubMed: 25956029). Activates the transcription of growth-related genes (PubMed: 24940000, PubMed: 25956029). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis

(PubMed:<u>24940000</u>, PubMed:<u>25956029</u>). Regulator of somatic

reprogramming, controls self-renewal of embryonic stem cells (By similarity).

Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production

of the PKM M2 isoform (PubMed:20010808).


Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome. Note=Association with chromatin is reduced by hyperphosphorylation (PubMed:30158517) Localization to the nucleolus is dependent on HEATR1 (PubMed:38225354)


Background

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'- CAC[GA]TG-3'. Activates the transcription of growth-related genes. Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed: 24940000).

Images



Western Blot analysis of various cells using Phospho-c-Myc (T58) Polyclonal Antibody diluted at 1:500



Western Blot analysis of 293 cells using Phospho-c-Myc (T58) Polyclonal Antibody diluted at 1:500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.