

CKR-5 (phospho Ser336) Polyclonal Antibody

Catalog # AP67221

Product Information

Application WB Primary Accession P51681

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 40524

Additional Information

Gene ID 1234

Other Names CCR5; CMKBR5; C-C chemokine receptor type 5; C-C CKR-5; CC-CKR-5; CCR-5;

CCR5; CHEMR13; HIV-1 fusion coreceptor; CD antigen CD195

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CCR5 (HGNC:1606)

Synonyms CMKBR5

Function Receptor for a number of inflammatory CC-chemokines including

CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. March 1997.

transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation. Participates in T-lymphocyte migration to the infection site by acting as a

chemotactic receptor (PubMed: 30713770).

Cell ular Location Cell membrane; Multi-pass membrane protein

Tissue Location Highly expressed in spleen, thymus, in the myeloid cell line THP-1, in the


promyeloblastic cell line KG-1a and on CD4+ and CD8+ T-cells. Medium levels in peripheral blood leukocytes and in small intestine. Low levels in ovary and

lung.

Background

Receptor for a number of inflammatory CC-chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation.

Images



Western Blot analysis of Jurkat cells using Phospho-CKR-5 (S336) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.