

Myf-5 (phospho Ser49) Polyclonal Antibody

Catalog # AP67327

Product Information

ApplicationWB, IHC-PPrimary AccessionP13349

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW28296

Additional Information

Gene ID 4617

Other Names MYF5; BHLHC2; Myogenic factor 5; Myf-5; Class C basic helix-loop-helix

protein 2; bHLHc2

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name MYF5

Synonyms BHLHC2

Function Transcriptional activator that promotes transcription of muscle-specific

target genes and plays a role in muscle differentiation (PubMed: 29887215). Together with MYOG and MYOD1, co-occupies muscle- specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into


myoblasts. Probable sequence specific DNA-binding protein.

Cellular Location Nucleus.

Background

Acts as a transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation. Together with MYOG and MYOD1, co-occupies muscle-specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. Probable sequence specific DNA-binding protein.

Images



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.