

TOMM70A Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6745b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q94826
Other Accession	Q75Q39 , Q9CZW5
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB20272
Antigen Region	579-608

Additional Information

Other Names	Mitochondrial import receptor subunit TOM70, Mitochondrial precursor proteins import receptor, Translocase of outer membrane 70 kDa subunit, TOMM70A, KIAA0719, TOM70
Target/Specificity	This TOMM70A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 579-608 amino acids from the C-terminal region of human TOMM70A.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TOMM70A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background

The translocase of outer mitochondrial membrane (TOM) complex is a multisubunit complex involved in the recognition, unfolding, and translocation of preproteins into the mitochondria.


References

Blesa,J.R., Gene 427 (1-2), 58-64 (2008)


Chou,C.H., Mol. Biol. Cell 17 (9), 3952-3963 (2006)

Edmonson,A.M., Cell Commun. Adhes. 9 (1), 15-27 (2002)


Images



Western blot analysis of TOMM70A antibody (C-term) (Cat. #AP6745b) in HL60 cell line lysates (35ug/lane). TOMM70A (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with TOMM70A Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of heLa cells using TOMM70A Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Citations

- [Proteomic analysis of mitochondria in respiratory epithelial cells infected with human respiratory syncytial virus and functional implications for virus and cell biology.](#)
- [Endothelial Mitochondrial Preprotein Translocase Tomm7-Rac1 Signaling Axis Dominates Cerebrovascular Network Homeostasis.](#)