

Nek9 (phospho Thr210) Polyclonal Antibody

Catalog # AP67496

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q8TD19
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	107168

Additional Information

Gene ID	91754
Other Names	NEK9; KIAA1995; NEK8; NERCC; Serine/threonine-protein kinase Nek9; Nercc1 kinase; Never in mitosis A-related kinase 9; NimA-related protein kinase 9; NimA-related kinase 8; Nek8
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	NEK9 {ECO:0000303 PubMed:12840024, ECO:0000312 HGNC:HGNC:18591}
Function	Pleiotropic regulator of mitotic progression, participating in the control of spindle dynamics and chromosome separation (PubMed: 12101123 , PubMed: 12840024 , PubMed: 14660563 , PubMed: 19941817). Phosphorylates different histones, myelin basic protein, beta-casein, and BICD2 (PubMed: 11864968). Phosphorylates histone H3 on serine and threonine residues and beta-casein on serine residues (PubMed: 11864968). Important for G1/S transition and S phase progression (PubMed: 12840024 , PubMed: 14660563 , PubMed: 19941817). Phosphorylates NEK6 and NEK7 and stimulates their activity by releasing the autoinhibitory functions of Tyr-108 and Tyr-97 respectively (PubMed: 12840024 , PubMed: 14660563 , PubMed: 19941817 , PubMed: 26522158).
Cellular Location	Cytoplasm. Nucleus
Tissue Location	Most abundant in heart, liver, kidney and testis. Also expressed in smooth

Background

Pleiotropic regulator of mitotic progression, participating in the control of spindle dynamics and chromosome separation. Phosphorylates different histones, myelin basic protein, beta-casein, and BICD2. Phosphorylates histone H3 on serine and threonine residues and beta-casein on serine residues. Important for G1/S transition and S phase progression. Phosphorylates NEK6 and NEK7 and stimulates their activity by releasing the autoinhibitory functions of Tyr-108 and Tyr-97 respectively.

Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.