

GABAA Rβ1 (phospho Ser434) Polyclonal Antibody

Catalog # AP67508

Product Information

Application WB, IHC-P **Primary Accession** P18505

Reactivity Human, Mouse, Rat, Monkey

HostRabbitClonalityPolyclonalCalculated MW54235

Additional Information

Gene ID 2560

Other Names GABRB1; Gamma-aminobutyric acid receptor subunit beta-1; GABA(A)

receptor subunit beta-1

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name GABRB1 (HGNC:4081)

Function Beta subunit of the heteropentameric ligand-gated chloride channel gated

by gamma-aminobutyric acid (GABA), a major inhibitory neurotransmitter in the brain (PubMed:10449790, PubMed:16412217, PubMed:26950270). GABA-gated chloride channels, also named GABA(A) receptors (GABAAR), consist of five subunits arranged around a central pore and contain one or two GABA active binding sites located at the alpha and beta subunit interfaces, depending on subunit composition (By similarity). When activated by GABA, GABAARs selectively allow the flow of chloride anions across the cell

membrane down their electrochemical gradient (PubMed: 10449790,

PubMed: 16412217, PubMed: 26950270). Chloride influx into the postsynaptic neuron following GABAAR opening decreases the neuron ability to generate a

new action potential, thereby reducing nerve transmission

(PubMed:<u>16412217</u>, PubMed:<u>26950270</u>). Beta-containing GABAARs can simultaneously bind GABA and histamine where histamine binds at the interface of two neighboring beta subunits, which may be involved in the


regulation of sleep and wakefulness (By similarity).

Postsynaptic cell membrane {ECO:0000250|UniProtKB:P08220}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P08220}. Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P08220}

Background

Component of the heteropentameric receptor for GABA, the major inhibitory neurotransmitter in the vertebrate brain. Functions also as histamine receptor and mediates cellular responses to histamine. Functions as receptor for diazepines and various anesthetics, such as pentobarbital; these are bound at a separate allosteric effector binding site. Functions as ligand- gated chloride channel.

Images



Western Blot analysis of various cells using Phospho-GABAA Rβ1 (S434) Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.