

# c-Myc (phospho Ser62) Polyclonal Antibody

Catalog # AP67793

## Product Information

---

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P01106</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50565

## Additional Information

---

Gene ID	4609
Other Names	MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	MYC
Synonyms	BHLHE39
Function	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed: <a href="#">24940000</a> , PubMed: <a href="#">25956029</a> ). Activates the transcription of growth-related genes (PubMed: <a href="#">24940000</a> , PubMed: <a href="#">25956029</a> ). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed: <a href="#">24940000</a> , PubMed: <a href="#">25956029</a> ). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed: <a href="#">20010808</a> ).


## Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome.  
Note=Association with chromatin is reduced by hyperphosphorylation  
(PubMed:30158517) Localization to the nucleolus is dependent on HEATR1  
(PubMed:38225354)


## Background

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'- CAC[GA]TG-3'. Activates the transcription of growth-related genes. Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:[24940000](#)).

## Images



Western Blot analysis of various cells using  
Phospho-c-Myc (S62) Polyclonal Antibody diluted at  
1 : 1000



Western Blot analysis of 293 cells using Phospho-c-Myc  
(S62) Polyclonal Antibody diluted at 1 : 1000

## Citations

- [Hyperthermia depletes Oct4 in mouse blastocysts and stem cells](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.