

# Rad51 (phospho Thr309) Polyclonal Antibody

Catalog # AP68038

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">Q06609</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	36966

## Additional Information

---

<b>Gene ID</b>	5888
<b>Other Names</b>	RAD51; RAD51A; RECA; DNA repair protein RAD51 homolog 1; HsRAD51; hRAD51; RAD51 homolog A
<b>Dilution</b>	WB~~Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	RAD51 ( <a href="#">HGNC:9817</a> )
<b>Synonyms</b>	RAD51A, RECA
<b>Function</b>	Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination (HR) (PubMed: <a href="#">12205100</a> , PubMed: <a href="#">18417535</a> , PubMed: <a href="#">20231364</a> , PubMed: <a href="#">20348101</a> , PubMed: <a href="#">22325354</a> , PubMed: <a href="#">23509288</a> , PubMed: <a href="#">23754376</a> , PubMed: <a href="#">26681308</a> , PubMed: <a href="#">28575658</a> , PubMed: <a href="#">32640219</a> ). Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed: <a href="#">12205100</a> , PubMed: <a href="#">18417535</a> , PubMed: <a href="#">20231364</a> , PubMed: <a href="#">20348101</a> , PubMed: <a href="#">23509288</a> , PubMed: <a href="#">23754376</a> , PubMed: <a href="#">26681308</a> , PubMed: <a href="#">28575658</a> ). Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template (PubMed: <a href="#">12205100</a> , PubMed: <a href="#">18417535</a> , PubMed: <a href="#">20231364</a> , PubMed: <a href="#">20348101</a> , PubMed: <a href="#">23509288</a> , PubMed: <a href="#">23754376</a> , PubMed: <a href="#">26681308</a> , PubMed: <a href="#">28575658</a> , PubMed: <a href="#">38459011</a> ). Recruited to

resolve stalled replication forks during replication stress (PubMed:[27797818](#), PubMed:[31844045](#)). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR (PubMed:[12442171](#), PubMed:[24141787](#)). Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3 (PubMed:[20413593](#)). Also involved in interstrand cross-link repair (PubMed:[26253028](#)).

## Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Mitochondrion matrix Chromosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Colocalizes with RAD51AP1 and RPA2 to multiple nuclear foci upon induction of DNA damage (PubMed:20154705). DNA damage induces an increase in nuclear levels (PubMed:20154705). Together with FIGNL1, redistributed in discrete nuclear DNA damage-induced foci after ionizing radiation (IR) or camptothecin (CPT) treatment (PubMed:23754376). Accumulated at sites of DNA damage in a SPIDR- dependent manner (PubMed:23509288). Recruited at sites of DNA damage in a MCM9-MCM8-dependent manner (PubMed:23401855). Recruited at sites of DNA damage following interaction with TOPBP1 in S-phase (PubMed:26811421). Colocalizes with ERCC5/XPG to nuclear foci in S phase (PubMed:26833090). Recruited to stalled replication forks during replication stress by the TONSL-MMS22L complex, as well as ATAD5 and WDR48 in an ATR-dependent manner (PubMed:27797818, PubMed:31844045)


## Tissue Location

Highly expressed in testis and thymus, followed by small intestine, placenta, colon, pancreas and ovary. Weakly expressed in breast

## Background

Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination (HR) (PubMed:[28575658](#)). Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template. Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:[26681308](#)). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3. Also involved in interstrand cross-link repair (PubMed:[26253028](#)).

## Images



Western blot analysis of HELA-UV 293T-UV using p-Rad51 (T309) antibody. Antibody was diluted at 1:500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).