

14-3-3 ζ Polyclonal Antibody

Catalog # AP68190

Product Information

Application	WB, IHC-P, IP, IF, ICC, E
Primary Accession	P63104
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27745

Additional Information

Gene ID	7534
Other Names	YWHAZ; 14-3-3 protein zeta/delta; Protein kinase C inhibitor protein 1; KCIP-1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A IP~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	YWHAZ
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: 14578935 , PubMed: 15071501 , PubMed: 15644438 , PubMed: 16376338 , PubMed: 16959763 , PubMed: 31024343 , PubMed: 9360956). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: 35662396). Binding generally results in the modulation of the activity of the binding partner (PubMed: 35662396). Promotes cytosolic retention and inactivation of TFEB transcription factor by binding to phosphorylated TFEB (PubMed: 35662396). Induces ARHGEF7 activity on RAC1 as well as lamellipodia and membrane ruffle formation (PubMed: 16959763). In neurons, regulates spine maturation through the modulation of ARHGEF7 activity (By similarity).
Cellular Location	Cytoplasm. Melanosome. Note=Located to stage I to stage IV melanosomes.

Background

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner.


Images



Western Blot analysis of various cells using 14-3-3 ζ Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of COLO205 cells using 14-3-3 ζ Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of 293T HELA using 14-3-3 ζ Polyclonal Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.