10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ABHD12 Polyclonal Antibody

Catalog # AP68238

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q8N2K0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45097

Additional Information

Gene ID	26090
Other Names	ABHD12; C20orf22; Monoacylglycerol lipase ABHD12; 2-arachidonoylglycerol hydrolase; Abhydrolase domain-containing protein 12
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	ABHD12 {ECO:0000303|PubMed:20797687, ECO:0000312|HGNC:HGNC:15868}
Function	Lysophosphatidylserine (LPS) lipase that mediates the hydrolysis of lysophosphatidylserine, a class of signaling lipids that regulates immunological and neurological processes (PubMed:25290914, PubMed:30237167, PubMed:30420694, PubMed:30643283, PubMed:30720278, PubMed:33571455). Represents a major lysophosphatidylserine lipase in the brain, thereby playing a key role in the central nervous system (By similarity). Prefers as substrates lysophosphatidylserines having long and very long chain acyl chains (PubMed:33571455). Also able to hydrolyze oxidized phosphatidylserine; oxidized phosphatidylserine is produced in response to severe inflammatory stress and constitutes a proapoptotic 'eat me' signal (PubMed:30643283). Also has monoacylglycerol (MAG) lipase activity: hydrolyzes 2- arachidonoylglycerol (2-AG), thereby acting as a regulator of endocannabinoid signaling pathways (PubMed:22969151, PubMed:24027063). Has a strong preference for

	very-long-chain lipid substrates; substrate specificity is likely due to improved catalysis and not improved substrate binding (PubMed:30237167).
Cellular Location	Endoplasmic reticulum membrane; Single-pass membrane protein


Background

Lysophosphatidylserine (LPS) lipase that plays a key role in the central nervous system. Represents a major LPS lipase in the brain (By similarity). May also have a 2- arachidonoylglycerol (2-AG) hydrolase activity and act as a regulator of endocannabinoid signaling pathways.

Images



Western Blot analysis of various cells using ABHD12 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human ovary. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.