10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

AI-BP Polyclonal Antibody

Catalog # AP68330

Product Information

Application	WB, IF, ICC, E
Primary Accession	Q8NCW5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31675

Additional Information

Gene ID	128240
Other Names	APOA1BP; AIBP; YJEFN1; NAD(P)H-hydrate epimerase; Apolipoprotein A-I-binding protein; AI-BP; NAD(P)HX epimerase; YjeF N-terminal domain-containing protein 1; YjeF_N1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	NAXE (HGNC:18453)
Function	Catalyzes the epimerization of the S- and R-forms of NAD(P)HX, a damaged form of NAD(P)H that is a result of enzymatic or heat-dependent hydration (By similarity) (PubMed:27616477). This is a prerequisite for the S-specific NAD(P)H-hydrate dehydratase to allow the repair of both epimers of NAD(P)HX (By similarity). Accelerates cholesterol efflux from endothelial cells to high-density lipoprotein (HDL) and thereby regulates angiogenesis (PubMed:23719382).
Cellular Location	Mitochondrion {ECO:0000255|HAMAP-Rule:MF_03159}. Secreted {ECO:0000255|HAMAP-Rule:MF_03159, ECO:0000269|PubMed:11991719}. Note=In sperm, secretion gradually increases during capacitation. {ECO:0000255|HAMAP-Rule:MF_03159}
Tissue Location	Ubiquitously expressed, with highest levels in kidney, heart and liver. Present in cerebrospinal fluid and urine but not in serum from healthy patients.

Present in serum of sepsis patients (at protein level).

Background

Catalyzes the epimerization of the S- and R-forms of NAD(P)HX, a damaged form of NAD(P)H that is a result of enzymatic or heat-dependent hydration. This is a prerequisite for the S- specific NAD(P)H-hydrate dehydratase to allow the repair of both epimers of NAD(P)HX.

Images



Western Blot analysis of various cells using AI-BP Polyclonal Antibody diluted at 1：2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.