

# AKAP 95 Polyclonal Antibody

Catalog # AP68351

## Product Information

---

Application	WB, IHC-P, IF
Primary Accession	<a href="#">O43823</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76108

## Additional Information

---

Gene ID	10270
Other Names	AKAP8; AKAP95; A-kinase anchor protein 8; AKAP-8; A-kinase anchor protein 95 kDa; AKAP 95
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	AKAP8
Synonyms	AKAP95
Function	Anchoring protein that mediates the subcellular compartmentation of cAMP-dependent protein kinase (PKA type II) (PubMed: <a href="#">9473338</a> ). Acts as an anchor for a PKA-signaling complex onto mitotic chromosomes, which is required for maintenance of chromosomes in a condensed form throughout mitosis. Recruits condensin complex subunit NCAPD2 to chromosomes required for chromatin condensation; the function appears to be independent from PKA-anchoring (PubMed: <a href="#">10601332</a> , PubMed: <a href="#">10791967</a> , PubMed: <a href="#">11964380</a> ). May help to deliver cyclin D/E to CDK4 to facilitate cell cycle progression (PubMed: <a href="#">14641107</a> ). Required for cell cycle G2/M transition and histone deacetylation during mitosis. In mitotic cells recruits HDAC3 to the vicinity of chromatin leading to deacetylation and subsequent phosphorylation at 'Ser-10' of histone H3; in this function may act redundantly with AKAP8L (PubMed: <a href="#">16980585</a> ). Involved in nuclear retention of RPS6KA1 upon ERK activation thus inducing cell proliferation

(PubMed:[22130794](#)). May be involved in regulation of DNA replication by acting as scaffold for MCM2 (PubMed:[12740381](#)). Enhances HMT activity of the KMT2 family MLL4/WBP7 complex and is involved in transcriptional regulation. In a teratocarcinoma cell line is involved in retinoic acid-mediated induction of developmental genes implicating H3 'Lys-4' methylation (PubMed:[23995757](#)). May be involved in recruitment of active CASP3 to the nucleus in apoptotic cells (PubMed:[16227597](#)). May act as a carrier protein of GJA1 for its transport to the nucleus (PubMed:[26880274](#)). May play a repressive role in the regulation of rDNA transcription. Preferentially binds GC-rich DNA in vitro. In cells, associates with ribosomal RNA (rRNA) chromatin, preferentially with rRNA promoter and transcribed regions (PubMed:[26683827](#)). Involved in modulation of Toll- like receptor signaling. Required for the cAMP-dependent suppression of TNF-alpha in early stages of LPS-induced macrophage activation; the function probably implicates targeting of PKA to NFKB1 (By similarity).

#### Cellular Location

Nucleus. Nucleus matrix. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q9DBR0}. Note=Associated with the nuclear matrix in interphase and redistributes mostly to chromatin at mitosis However, mitotic chromatin localization has been questioned. Upon nuclear reassembly at the end of mitosis, is sequestered into the daughter nuclei where it re-acquires an interphase distribution Exhibits partial localization to the nucleolus in interphase, where it colocalizes with UBTF/UBF, suggesting localization to the fibrillary center and/or to the dense fibrillary component. Colocalizes with GJA1 at the nuclear membrane specifically during cell cycle G1/S phase

#### Tissue Location


Highly expressed in heart, liver, skeletal muscle, kidney and pancreas. Expressed in mature dendritic cells

## Background


Anchoring protein that mediates the subcellular compartmentation of cAMP-dependent protein kinase (PKA type II) (PubMed:[9473338](#)). Acts as an anchor for a PKA-signaling complex onto mitotic chromosomes, which is required for maintenance of chromosomes in a condensed form throughout mitosis. Recruits condensin complex subunit NCAPD2 to chromosomes required for chromatin condensation; the function appears to be independent from PKA-anchoring (PubMed:[10601332](#), PubMed:[10791967](#), PubMed:[11964380](#)). May help to deliver cyclin D/E to CDK4 to facilitate cell cycle progression (PubMed:[14641107](#)). Required for cell cycle G2/M transition and histone deacetylation during mitosis. In mitotic cells recruits HDAC3 to the vicinity of chromatin leading to deacetylation and subsequent phosphorylation at 'Ser-10' of histone H3; in this function may act redundantly with AKAP8L (PubMed:[16980585](#)). Involved in nuclear retention of RPS6KA1 upon ERK activation thus inducing cell proliferation (PubMed:[22130794](#)). May be involved in regulation of DNA replication by acting as scaffold for MCM2 (PubMed:[12740381](#)). Enhances HMT activity of the KMT2 family MLL4/WBP7 complex and is involved in transcriptional regulation. In a teratocarcinoma cell line is involved in retinoic acid-mediated induction of developmental genes implicating H3 'Lys-4' methylation (PubMed:[23995757](#)). May be involved in recruitment of active CASP3 to the nucleus in apoptotic cells (PubMed:[16227597](#)). May act as a carrier protein of GJA1 for its transport to the nucleus (PubMed:[26880274](#)). Seems to involved in modulation of rDNA transcription. Preferentially binds GC-rich DNA in vitro and associates to GC-rich ribosomal RNA promoters (PubMed:[26683827](#)). Involved in modulation of Toll-like receptor signaling. Required for the cAMP-dependent suppression of TNF-alpha in early stages of LPS-induced macrophage activation; the function probably implicates targeting of PKA to NFKB1 (By similarity).

## Images

Western Blot analysis of various cells using AKAP 95 Polyclonal Antibody diluted at 1 : 1000 cells nucleus



extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of MCF7 cells using AKAP 95 Polyclonal Antibody diluted at 1 : 1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.