

# DFNA5 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6842A

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">O60443</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB20936
Calculated MW	54555
Antigen Region	59-87

## Additional Information

---

Gene ID	1687
Other Names	Non-syndromic hearing impairment protein 5, Inversely correlated with estrogen receptor expression 1, ICERE-1, DFNA5, ICERE1
Target/Specificity	This DFNA5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 59-87 amino acids from the N-terminal region of human DFNA5.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DFNA5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	GSDME {ECO:0000303   PubMed:28459430, ECO:0000312   HGNC:HGNC:2810}
Function	[Gasdermin-E]: Precursor of a pore-forming protein that converts non-inflammatory apoptosis to pyroptosis (PubMed: <a href="#">27281216</a> , PubMed: <a href="#">28459430</a> , PubMed: <a href="#">33852854</a> , PubMed: <a href="#">35594856</a> , PubMed: <a href="#">36607699</a> ). This form constitutes the precursor of the pore-forming

protein: upon cleavage, the released N-terminal moiety (Gasdermin-E, N-terminal) binds to membranes and forms pores, triggering pyroptosis (PubMed:[28459430](#)).

#### Cellular Location

[Gasdermin-E, N-terminal]: Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q5Y4Y6}

#### Tissue Location

Expressed in cochlea (PubMed:9771715). Low level of expression in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas, with highest expression in placenta (PubMed:9771715).

## Background

---

Hearing impairment is a heterogeneous condition with over 40 loci described. DFNA5 is expressed in fetal cochlea, however, its function is not known.


## References

---


Cheng,J., et.al., Clin. Genet. 72 (5), 471-477 (2007)

## Images

---




Western blot analysis of DFNA5 Antibody (N-term) (Cat. #AP6842a) in HeLa cell line lysates (35ug/lane). DFNA5 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human breast carcinoma reacted with DFNA5 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Flow cytometric analysis of MDA-MB468 cells using DFNA5 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram)FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.