

API5 Polyclonal Antibody

Catalog # AP68447

Product Information

Application WB, IHC-P, IF, ICC, E

Primary Accession Q9BZZ5

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW59005

Additional Information

Gene ID 8539

Other Names API5; MIG8; Apoptosis inhibitor 5; API-5; Antiapoptosis clone 11 protein;

AAC-11; Cell migration-inducing gene 8 protein; Fibroblast growth factor

2-interacting factor; FIF; Protein XAGL

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A

IF~~1:50~200 ICC~~N/A E~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name API5 (HGNC:594)

Function Antiapoptotic factor that may have a role in protein assembly. Negatively

regulates ACIN1. By binding to ACIN1, it suppresses ACIN1 cleavage from CASP3 and ACIN1-mediated DNA fragmentation. Also known to efficiently suppress E2F1-induced apoptosis. Its depletion enhances the cytotoxic action

of the chemotherapeutic drugs.

Cellular Location Nucleus. Cytoplasm. Note=Mainly nuclear. Can also be cytoplasmic

Tissue Location Expressed in all tissues tested, including heart, brain, placenta, lung, liver,

skeletal muscle, kidney and pancreas Highest levels in heart, pancreas and placenta. Highly expressed in several cancers. Preferentially expressed in squamous cell carcinoma versus adenocarcinoma in non-small cell lung


cancer

Background

Antiapoptotic factor that may have a role in protein assembly. Negatively regulates ACIN1. By binding to ACIN1, it suppresses ACIN1 cleavage from CASP3 and ACIN1-mediated DNA fragmentation. Also known to efficiently suppress E2F1-induced apoptosis. Its depletion enhances the cytotoxic action of the chemotherapeutic drugs.

Images

19-



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.