

# Arp3 Polyclonal Antibody

Catalog # AP68519

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P61158</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	47371

## Additional Information

---

<b>Gene ID</b>	10096
<b>Other Names</b>	ACTR3; ARP3; Actin-related protein 3; Actin-like protein 3
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---


<b>Name</b>	ACTR3
<b>Synonyms</b>	ARP3
<b>Function</b>	ATP-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF) (PubMed: <a href="#">9000076</a> ). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility (PubMed: <a href="#">9000076</a> ). Seems to contact the pointed end of the daughter actin filament (PubMed: <a href="#">9000076</a> ). In podocytes, required for the formation of lamellipodia downstream of AVIL and PLCE1 regulation (PubMed: <a href="#">29058690</a> ). In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed: <a href="#">17220302</a> , PubMed: <a href="#">29925947</a> ). The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed: <a href="#">29925947</a> ). Plays a role in ciliogenesis (PubMed: <a href="#">20393563</a> ).
<b>Cellular Location</b>	Cytoplasm, cytoskeleton. Cell projection. Nucleus. Note=In pre- apoptotic

cells, colocalizes with MEFV in large specks (pyroptosomes)  
(PubMed:19109554)

## Background

ATP-binding component of the Arp2/3 complex, a multiprotein complex that mediates actin polymerization upon stimulation by nucleation-promoting factor (NPF) (PubMed:[9000076](#)). The Arp2/3 complex mediates the formation of branched actin networks in the cytoplasm, providing the force for cell motility (PubMed:[9000076](#)). Seems to contact the pointed end of the daughter actin filament (PubMed:[9000076](#)). In addition to its role in the cytoplasmic cytoskeleton, the Arp2/3 complex also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:[17220302](#), PubMed:[29925947](#)). The Arp2/3 complex promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double- strand breaks (DSBs) (PubMed:[29925947](#)). Plays a role in ciliogenesis (PubMed:[20393563](#)).

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.