

BMP-5 Polyclonal Antibody

Catalog # AP68684

Product Information

Application WB, IHC-P **Primary Accession** P22003

Reactivity Human, Mouse

HostRabbitClonalityPolyclonalCalculated MW51737

Additional Information

Gene ID 653

Other Names BMP5; Bone morphogenetic protein 5; BMP-5

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name BMP5

Function Growth factor of the TGF-beta superfamily that plays essential roles in many

developmental processes, including cartilage and bone formation or

neurogenesis (PubMed: 11580864, PubMed: 29321139). Initiates the canonical BMP signaling cascade by associating with type I receptor BMPR1A and type II receptor BMPR2 (PubMed: 11580864). In turn, BMPR1A propagates signal by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressers of transcription of target genes (PubMed: 11580864).

and repressors of transcription of target genes (PubMed: 11580864,

PubMed:<u>29321139</u>). Can also signal through non-canonical pathway such as MAPK p38 signaling cascade to promote chondrogenic differentiation (PubMed:<u>20402566</u>). Promotes the expression of HAMP, this is repressed by

its interaction with ERFE (PubMed:30097509).


Cellular Location Secreted.

Tissue Location Expressed in the lung and liver.


Background

Induces cartilage and bone formation.

Images



Western Blot analysis of various cells using BMP-5 Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HeLa cells using BMP-5 Polyclonal Antibody diluted at 1: 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.