

C1INH Polyclonal Antibody

Catalog # AP68744

Product Information

Application	WB, IHC-P
Primary Accession	<u>P05155</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55154

Additional Information

Gene ID	710
Other Names	SERPING1; C1IN; C1NH; Plasma protease C1 inhibitor; C1 Inh; C1Inh; C1 esterase inhibitor; C1-inhibiting factor; Serpin G1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	SERPING1
Synonyms	C1IN, C1NH
Function	Serine protease inhibitor, which acrs as a regulator of the classical complement pathway (PubMed: <u>10946292</u> , PubMed: <u>11527969</u> , PubMed: <u>3458172</u> , PubMed: <u>6416294</u>). Forms a proteolytically inactive stoichiometric complex with the C1r or C1s proteases (PubMed: <u>10946292</u> , PubMed: <u>3458172</u> , PubMed: <u>6416294</u>). May also regulate blood coagulation, fibrinolysis and the generation of kinins (PubMed: <u>8495195</u>). Very efficient inhibitor of FXIIa. Inhibits chymotrypsin and kallikrein (PubMed: <u>8495195</u>).
Cellular Location	Secreted

Background

Activation of the C1 complex is under control of the C1- inhibitor. It forms a proteolytically inactive stoichiometric complex with the C1r or C1s proteases. May play a potentially crucial role in regulating

important physiological pathways including complement activation, blood coagulation, fibrinolysis and the generation of kinins. Very efficient inhibitor of FXIIa. Inhibits chymotrypsin and kallikrein.





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.