

# Carbonyl Reductase 3 Polyclonal Antibody

Catalog # AP68816

## Product Information

---

Application	WB, IHC-P
Primary Accession	<a href="#">O75828</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30850

## Additional Information

---

Gene ID	874
Other Names	CBR3; Carbonyl reductase [NADPH] 3; NADPH-dependent carbonyl reductase 3
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

---

Name	CBR3 ( <a href="#">HGNC:1549</a> )
Function	Catalyzes the NADPH-dependent reduction of carbonyl compounds to their corresponding alcohols (PubMed: <a href="#">18493841</a> ). Has low NADPH- dependent oxidoreductase activity. Acts on several orthoquinones, acts as well on non-quinone compounds, such as isatin or on the anticancer drug oracin (PubMed: <a href="#">15537833</a> , PubMed: <a href="#">18493841</a> , PubMed: <a href="#">19841672</a> ). Best substrates for CBR3 is 1,2- naphthoquinone, hence could play a role in protection against cytotoxicity of exogenous quinones (PubMed: <a href="#">19841672</a> ). Exerts activity toward ortho-quinones but not paraquinones. No endogenous substrate for CBR3 except isatin has been identified (PubMed: <a href="#">19841672</a> ).
Cellular Location	Cytoplasm.
Tissue Location	Detected in ovary, pancreas, intestine, colon, kidney, brain, thymus, lung, heart, liver, spleen, leukocyte, prostate and testis.

## Background

---

Has low NADPH-dependent oxidoreductase activity towards 4-benzoylpyridine and menadione (in vitro).

## Images

---



Western Blot analysis of various cells using Carbonyl Reductase 3 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.