

# HADHA Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6882b

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P40939</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	83000
<b>Antigen Region</b>	737-763

## Additional Information

---

<b>Gene ID</b>	3030
<b>Other Names</b>	Trifunctional enzyme subunit alpha, mitochondrial, 78 kDa gastrin-binding protein, TP-alpha, Long-chain enoyl-CoA hydratase, Long chain 3-hydroxyacyl-CoA dehydrogenase, HADHA, HADH
<b>Target/Specificity</b>	This HADHA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 737-763 amino acids from the C-terminal region of human HADHA.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	HADHA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	HADHA
<b>Synonyms</b>	HADH
<b>Function</b>	Mitochondrial trifunctional enzyme catalyzes the last three of the four

reactions of the mitochondrial beta-oxidation pathway (PubMed:[1550553](#), PubMed:[29915090](#), PubMed:[30850536](#), PubMed:[8135828](#), PubMed:[31604922](#)). The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA (PubMed:[29915090](#)). Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids (PubMed:[30850536](#), PubMed:[31604922](#)). Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA described here carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities while the trifunctional enzyme subunit beta/HADHB bears the 3-ketoacyl-CoA thiolase activity (PubMed:[29915090](#), PubMed:[30850536](#), PubMed:[8135828](#)). Independently of subunit beta, HADHA also exhibits a cardiolipin acyltransferase activity that participates in cardiolipin remodeling; cardiolipin is a major mitochondrial membrane phospholipid (PubMed:[23152787](#), PubMed:[31604922](#)). HADHA may act downstream of Tafazzin/TAZ, that remodels monolysocardiolipin (MLCL) to a cardiolipin intermediate, and then HADHA may continue to remodel this species into mature tetralinoleoyl-cardiolipin (PubMed:[31604922](#)). Has also been proposed to act directly on MLCL; capable of acylating MLCL using different acyl-CoA substrates, with highest activity for oleoyl-CoA (PubMed:[23152787](#)).

## Cellular Location

Mitochondrion. Mitochondrion inner membrane Note=Protein stability and association with mitochondrion inner membrane do not require HADHB.


## Background

HADHA is the alpha subunit of the mitochondrial trifunctional protein, which catalyzes the last three steps of mitochondrial beta-oxidation of long chain fatty acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four beta subunits, with the alpha subunit catalyzing the 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratase activities.

## References


Sims,H.F., et.al., Proc. Natl. Acad. Sci. U.S.A. 92 (3), 841-845 (1995)

## Images




All lanes: Anti-HADHA Antibody (C-term) at 1:2000  
 Lane 1: HeLa whole cell lysate  
 Lane 2: 293T whole cell lysate  
 Lane 3: A431 whole cell lysate  
 Lane 4: K562 whole cell lysate  
 Lane 5: Mouse heart lysate  
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 83KDa  
 Blocking/Dilution buffer: 5% NFDM/TBST.


All lanes : Anti-HADHA Antibody (C-term) at 1:1000  
 Lane 1: HeLa whole cell lysate  
 Lane 2: Jurkat whole cell lysate  
 Lysates/proteins at 20 µg per lane.  
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase




conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



HADHA Antibody (C-term) (Cat. #AP6882b) western blot analysis in A431 cell line lysates (35ug/lane). This demonstrates the HADHA antibody detected the HADHA protein (arrow).



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with HADHA Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



HADHA Antibody (C-term) (Cat. #AP6882b) flow cytometry analysis of Ramos cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.