

CCK-BR Polyclonal Antibody

Catalog # AP68891

Product Information

Application	WB, E
Primary Accession	P32239
Reactivity	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48419

Additional Information

Gene ID	887
Other Names	CCKBR; CCKRB; Gastrin/cholecystokinin type B receptor; CCK-B receptor; CCK-BR; Cholecystokinin-2 receptor; CCK2-R
Dilution	WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	CCKBR (HGNC:1571)
Synonyms	CCKRB
Function	Receptor for the peptide hormones gastrin and cholecystokinin (CCK). Expressed throughout the central nervous system, where it modulates processes such as anxiety, analgesia, arousal and neuroleptic activity. Couples to both GNA1 and GNAQ signaling pathways, but not to GNAS (PubMed: 34556863). Upon gastrin activation, reduces glucose absorption in intestinal epithelial cells by downregulating SGLT1 and GLUT2 expression through suppression of the PI3K/Akt/eIF4B pathway (By similarity). In the kidney, decreases SGLT2 expression under high- glucose conditions via ERK/NF-kappa-B signaling (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein.
Tissue Location	Isoform 1 is expressed in brain, pancreas, stomach, the colon cancer cell line LoVo and the T-lymphoblastoma Jurkat, but not in heart, placenta, liver, lung, skeletal muscle, kidney or the stomach cancer cell line AGS. Expressed at high

levels in the small cell lung cancer cell line NCI-H510, at lower levels in NCI-H345, NCI- H69 and GLC-28 cell lines, not expressed in GLC-19 cell line. Within the stomach, expressed at high levels in the mucosa of the gastric fundus and at low levels in the antrum and duodenum. Isoform 2 is present in pancreatic cancer cells and colorectal cancer cells, but not in normal pancreas or colonic mucosa. Isoform 3 is expressed in brain, pancreas, stomach, the stomach cancer cell line AGS and the colon cancer cell line LoVo.

Background

Receptor for gastrin and cholecystokinin. The CCK-B receptors occur throughout the central nervous system where they modulate anxiety, analgesia, arousal, and neuroleptic activity. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system.

Images



Western Blot analysis of various cells using CCK-BR Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.