

# CD137L Polyclonal Antibody

Catalog # AP68906

## Product Information

---

<b>Application</b>	WB, IF
<b>Primary Accession</b>	<a href="#">P41273</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	26625

## Additional Information

---

<b>Gene ID</b>	8744
<b>Other Names</b>	TNFSF9; Tumor necrosis factor ligand superfamily member 9; 4-1BB ligand; 4-1BBL
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information


---

<b>Name</b>	TNFSF9
<b>Function</b>	Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation- induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.
<b>Cellular Location</b>	Membrane; Single-pass type II membrane protein.
<b>Tissue Location</b>	Expressed in brain, placenta, lung, skeletal muscle and kidney


## Background

---

Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.



Western Blot analysis of various cells using CD137L Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HuvEc cells using CD137L Polyclonal Antibody diluted at 1 : 1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.