

CD137L Polyclonal Antibody

Catalog # AP68906

Product Information

ApplicationWB, IFPrimary AccessionP41273ReactivityHumanHostRabbitClonalityPolyclonalCalculated MW26625

Additional Information

Gene ID 8744

Other Names TNFSF9; Tumor necrosis factor ligand superfamily member 9; 4-1BB ligand;

4-1BBL

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000.

ELISA: 1/20000. Not yet tested in other applications. IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name TNFSF9

Function Cytokine that binds to TNFRSF9. Induces the proliferation of activated

peripheral blood T-cells. May have a role in activation- induced cell death (AICD). May play a role in cognate interactions between T-cells and

B-cells/macrophages.


Cellular Location Membrane; Single-pass type II membrane protein.

Tissue Location Expressed in brain, placenta, lung, skeletal muscle and kidney


Background

Cytokine that binds to TNFRSF9. Induces the proliferation of activated peripheral blood T-cells. May have a role in activation-induced cell death (AICD). May play a role in cognate interactions between T-cells and B-cells/macrophages.

Images



Western Blot analysis of various cells using CD137L Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of HuvEc cells using CD137L Polyclonal Antibody diluted at 1:1000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.