

CD83 Polyclonal Antibody

Catalog # AP68962

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q01151
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	23042

Additional Information

Gene ID	9308
Other Names	CD83; CD83 antigen; hCD83; B-cell activation protein; Cell surface protein HB15; CD antigen CD83
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C


Protein Information

Name	CD83
Function	Transmembrane glycoprotein predominantly found on the surface of many immune cells including dendritic cells or lymphocytes that plays various roles in immune response regulation. Plays an essential role in CD4(+) T-selection, differentiation and stability by regulating the activity of the major E3 ubiquitin ligase responsible for controlling MHCII trafficking MARCHF8. Also inhibits MARCHF1 association with MHCII or CD86 to prevent their ubiquitination and subsequent degradation (PubMed: 21220452). In addition, acts as an important modulator of protective responses against acute infections (By similarity).
Cellular Location	Membrane; Single-pass type I membrane protein.
Tissue Location	Expressed by activated lymphocytes, Langerhans cells and activated dendritic cells.


Background

May play a significant role in antigen presentation or the cellular interactions that follow lymphocyte activation.


Images




Western Blot analysis of various cells using CD83 Polyclonal Antibody diluted at 1 : 500



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).


Immunohistochemical analysis of paraffin-embedded Human stomach. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human stomach. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human stomach. 1, Antibody was diluted at 1:400(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.