10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Cdc27 Polyclonal Antibody

Catalog # AP68985

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	P30260
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	91867

Additional Information

Gene ID	996
Other Names	CDC27; ANAPC3; D0S1430E; D17S978E; Cell division cycle protein 27 homolog; Anaphase-promoting complex subunit 3; APC3; CDC27 homolog; CDC27Hs; H-NUC
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20℃

Protein Information


Name	CDC27
Synonyms	ANAPC3, D0S1430E, D17S978E
Function	Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through the cell cycle (PubMed:18485873, PubMed:27120157, PubMed:27509861). APC/C acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:18485873). APC/C catalyzes assembly of branched 'Lys-11'-/'Lys- 48'-linked branched ubiquitin chains on target proteins (PubMed:29033132). APC/C is activated by CDC20 in the metaphase/anaphase transition of cell cycle, targeting the degradation of cyclin B and securin (PubMed:27120157, PubMed:27509861). APC/C is regulated by the mitotic checkpoint complex (MCC), which inhibits APC/C and delays chromosome segregation (PubMed:27509861).

Cellular Location Nucleus. Cytoplasm, cytoskeleton, spindle

Background

Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains.

Images



Western Blot analysis of various cells using Cdc27 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.