

Cdc27 Polyclonal Antibody

Catalog # AP68985

Product Information

Application WB, IHC-P Primary Accession P30260

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW91867

Additional Information

Gene ID 996

Other Names CDC27; ANAPC3; D0S1430E; D17S978E; Cell division cycle protein 27 homolog;

Anaphase-promoting complex subunit 3; APC3; CDC27 homolog; CDC27Hs;

H-NUC

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CDC27

Synonyms ANAPC3, D0S1430E, D17S978E

Function Component of the anaphase promoting complex/cyclosome (APC/C), a cell

cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle (PubMed:18485873). The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (PubMed:18485873). The APC/C complex catalyzes assembly of branched 'Lys-11'-/'Lys-48'-linked branched ubiquitin chains on


target proteins (PubMed:29033132).

Cellular Location Nucleus. Cytoplasm, cytoskeleton, spindle

Background

Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.