

CDK5 Activator-binding C42 Polyclonal Antibody

Catalog # AP69019

Product Information

Application	IHC-P
Primary Accession	Q96SZ6
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67689

Additional Information

Gene ID	51654
Other Names	CDK5RAP1; C20orf34; CGI-05; HSPC167; CDK5 regulatory subunit-associated protein 1; CDK5 activator-binding protein C42
Dilution	IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information


Name	CDK5RAP1 (HGNC:15880)
Synonyms	C20orf34
Function	Methylthiotransferase that catalyzes the conversion of N6-(dimethylallyl)adenosine (i(6)A) to 2-methylthio-N6-(dimethylallyl)adenosine (ms(2)i(6)A) at position 37 (adjacent to the 3'-end of the anticodon) of four mitochondrial DNA-encoded tRNAs (Ser(UCN), Phe, Tyr and Trp) (PubMed: 22422838 , PubMed: 25738458 , PubMed: 28981754). Essential for efficient and highly accurate protein translation by the ribosome (PubMed: 22422838 , PubMed: 25738458 , PubMed: 28981754). Specifically inhibits CDK5 activation by CDK5R1 (PubMed: 11882646). Essential for efficient mitochondrial protein synthesis and respiratory chain; shows pathological consequences in mitochondrial disease (PubMed: 25738458).
Cellular Location	Mitochondrion.
Tissue Location	Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas (PubMed:10721722). Expressed in neurons of central nervous tissue (PubMed:10721722, PubMed:11882646) [Isoform 6]: High expression in

placenta and lung.

Background

Specifically inhibits CDK5 activation by CDK5R1.

Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.