

CEP78 Polyclonal Antibody

Catalog # AP69049

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q5JTW2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76396

Additional Information

Gene ID	84131
Other Names	CEP78; C9orf81; Centrosomal protein of 78 kDa; Cep78
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CEP78 {ECO:0000303 PubMed:27588451, ECO:0000312 HGNC:HGNC:25740}
Function	Centriole wall protein that localizes to mature centrioles and regulates centriole and cilia biogenesis (PubMed: 27246242 , PubMed: 27588451 , PubMed: 28242748 , PubMed: 34259627). Involved in centrosome duplication: required for efficient PLK4 centrosomal localization and PLK4-induced overduplication of centrioles (PubMed: 27246242). Involved in cilium biogenesis and controls cilium length (PubMed: 27588451). Acts as a regulator of protein stability by preventing ubiquitination of centrosomal proteins, such as CCP110 and tektins (PubMed: 28242748 , PubMed: 34259627). Associates with the EDVP complex, preventing ubiquitination and degradation of CCP110 (PubMed: 28242748 , PubMed: 34259627). Promotes deubiquitination of tektin proteins (TEKT1, TEKT2, TEK3, TEKT4 and TEKT5) via its interaction with USP16 (By similarity).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly localizes at

the centriolar wall, but also found in the pericentriolar material (PubMed:27246242). Expressed in photoreceptor inner segment (PubMed:27588452).


Tissue Location

Widely expressed (PubMed:27588451, PubMed:27588452). Expressed in different retinal cell types with higher expression in cone compared to rod cells (at protein level) (PubMed:27588452).

Background

May be required for efficient PLK4 centrosomal localization and PLK4-induced overduplication of centrioles (PubMed:[27246242](#)). May play a role in cilium biogenesis (PubMed:[27588451](#)).

Images



Western Blot analysis of various cells using CEP78 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.