

# Claudin-4 Polyclonal Antibody

Catalog # AP69133

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">O14493</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22077

## Additional Information

Gene ID	1364
Other Names	CLDN4; CPER; CPETR1; WBSCR8; Claudin-4; Clostridium perfringens enterotoxin receptor; CPE-R; CPE-receptor; Williams-Beuren syndrome chromosomal region 8 protein
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information


Name	CLDN4 {ECO:0000303   PubMed:35773259, ECO:0000312   HGNC:HGNC:2046}
Function	Can associate with other claudins to regulate tight junction structural and functional strand dynamics (PubMed: <a href="#">35773259</a> , PubMed: <a href="#">36008380</a> ). May coassemble with CLDN8 into tight junction strands containing anion-selective channels that convey paracellular chloride permeability in renal collecting ducts (By similarity) (PubMed: <a href="#">36008380</a> ). May integrate into CLDN3 strands to modulate localized tight junction barrier properties (PubMed: <a href="#">35773259</a> , PubMed: <a href="#">36008380</a> ). May disrupt strand assembly of channel-forming CLDN2 and CLDN15 and inhibit cation conductance (PubMed: <a href="#">35773259</a> , PubMed: <a href="#">36008380</a> ). Cannot form tight junction strands on its own (PubMed: <a href="#">35773259</a> , PubMed: <a href="#">36008380</a> ).
Cellular Location	Cell junction, tight junction. Cell membrane; Multi-pass membrane protein

## Background

Channel-forming tight junction protein that mediates paracellular chloride transport in the kidney. Plays a critical role in the paracellular reabsorption of filtered chloride in the kidney collecting ducts. Claudins play a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity.

## Images

---



Western Blot analysis of mouse-BRAIN cells using Claudin-4 Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.