

CNOT7 Polyclonal Antibody

Catalog # AP69185

Product Information

Application	WB, IHC-P, IF, ICC, E
Primary Accession	Q9UIV1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32745

Additional Information

Gene ID	29883
Other Names	CNOT7; CAF1; CCR4-NOT transcription complex subunit 7; BTG1-binding factor 1; CCR4-associated factor 1; CAF-1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CNOT7
Synonyms	CAF1
Function	Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate (PubMed: 19276069 , PubMed: 20634287 , PubMed: 31439799). Its function seems to be partially redundant with that of CNOT8 (PubMed: 19605561). Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation (PubMed: 19276069 , PubMed: 20634287 , PubMed: 31439799). During miRNA-mediated repression the complex also seems to act as translational repressor during translational initiation (PubMed: 20065043). Additional complex functions may be a consequence of its influence on mRNA expression (PubMed: 19276069 , PubMed: 23236473). Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti- proliferative activity (PubMed: 19276069 , PubMed: 23236473).


Cellular Location

Nucleus. Cytoplasm, P-body {ECO:0000250|UniProtKB:Q60809}. Cytoplasm, Cytoplasmic ribonucleoprotein granule. Note=NANOS2 promotes its localization to P-body (By similarity). Recruited to cytoplasmic ribonucleoprotein membraneless compartments by CAPRIN1, promoting deadenylation of mRNAs (PubMed:31439799) {ECO:0000250|UniProtKB:Q60809, ECO:0000269|PubMed:31439799}

Background

Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate. Its function seems to be partially redundant with that of CNOT8. Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. During miRNA-mediated repression the complex seems also to act as translational repressor during translational initiation. Additional complex functions may be a consequence of its influence on mRNA expression. Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti- proliferative activity.

Images



Western Blot analysis of various cells using CNOT7 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.