

# **CREB-1 Polyclonal Antibody**

Catalog # AP69284

### **Product Information**

Application WB, IHC-P Primary Accession P16220

**Reactivity** Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW35136

## **Additional Information**

**Gene ID** 1385

Other Names CREB1; Cyclic AMP-responsive element-binding protein 1; CREB-1;

cAMP-responsive element-binding protein 1

**Dilution** WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

#### **Protein Information**

Name CREB1

**Function** Phosphorylation-dependent transcription factor that stimulates

transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters (By similarity). Transcription activation is enhanced by the TORC coactivators which act independently of Ser-119 phosphorylation (PubMed:14536081). Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells (By similarity). Regulates

the expression of apoptotic and inflammatory response factors in

cardiomyocytes in response to ERFE-mediated activation of AKT signaling (By

similarity).


**Cellular Location** Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00312,

ECO:0000255 | PROSITE-ProRule:PRU00978, ECO:0000269 | PubMed:12552083}

# **Background**

Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells.

# **Images**



Western Blot analysis of various cells using CREB-1 Polyclonal Antibody diluted at 1 : 500

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.