

CRF-RII Polyclonal Antibody

Catalog # AP69293

Product Information

Application WB, IHC-P, IF **Primary Accession** Q13324

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 47688

Additional Information

Gene ID 1395

Other Names CRHR2; CRF2R; CRH2R; Corticotropin-releasing factor receptor 2; CRF-R-2;

CRF-R2; CRFR-2; Corticotropin-releasing hormone receptor 2; CRH-R-2; CRH-R2

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other

applications. IHC-P~~N/A IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CRHR2

Synonyms CRF2R, CRH2R

Function G-protein coupled receptor for CRH (corticotropin-releasing factor), UCN

(urocortin), UCN2 and UCN3. Has high affinity for UCN. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and down-stream effectors, such as adenylate cyclase.

Promotes the activation of adenylate cyclase, leading to increased

intracellular cAMP levels.


Cellular Location Cell membrane; Multi-pass membrane protein

Background

G-protein coupled receptor for CRH (corticotropin- releasing factor), UCN (urocortin), UCN2 and UCN3. Has high affinity for UCN. Ligand binding causes a conformation change that triggers signaling via guanine

nucleotide-binding proteins (G proteins) and down-stream effectors, such as adenylate cyclase. Promotes the activation of adenylate cyclase, leading to increased intracellular cAMP levels.

Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.