

# TPH1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6930a

## Product Information

---

|                          |   |
|--------------------------|---|
| <b>Application</b>       | IHC-P, FC, IF, WB, E  |
| <b>Primary Accession</b> | <a href="#">P17752</a>  |
| <b>Other Accession</b>   | <a href="#">Q92142</a> , <a href="#">P09810</a> , <a href="#">P17290</a> , <a href="#">P70080</a> |
| <b>Reactivity</b>        | Human   |
| <b>Predicted</b>         | Chicken, Rabbit, Rat, Xenopus   |
| <b>Host</b>              | Rabbit  |
| <b>Clonality</b>         | Polyclonal  |
| <b>Isotype</b>           | Rabbit IgG  |
| <b>Clone Names</b>       | RB21510   |
| <b>Calculated MW</b>     | 50985   |
| <b>Antigen Region</b>    | 35-62   |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 7166   |
| <b>Other Names</b>        | Tryptophan 5-hydroxylase 1, Tryptophan 5-monooxygenase 1, TPH1, TPH, TPRH, TRPH  |
| <b>Target/Specificity</b> | This TPH1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 35-62 amino acids from the N-terminal region of human TPH1.           |
| <b>Dilution</b>           | IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.  |
| <b>Format</b>             | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                      |
| <b>Precautions</b>        | TPH1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.   |

## Protein Information

---

|                 |                 |
|-----------------|-----------------|
| <b>Name</b>     | TPH1            |
| <b>Synonyms</b> | TPH, TPRH, TRPH |

|                        |  |
|------------------------|--|
| <b>Function</b>        | Oxidizes L-tryptophan to 5-hydroxy-L-tryptophan in the rate- determining step of serotonin biosynthesis. |
| <b>Tissue Location</b> | [Isoform 2]: Seems to be less widely expressed than isoform 1.   |


## Background

TPH1 is a member of the aromatic amino acid hydroxylase family. This protein catalyzes the first and rate limiting step in the biosynthesis of serotonin, an important hormone and neurotransmitter.


## References

Johansson,T.A., et.al., BMC Cancer 9, 321 (2009)  
Kuhn,D.M., et.al., J. Neurochem. 68 (5), 2220-2223 (1997)


## Images




Anti-TPH1 Antibody (N-term) at 1:500 dilution + 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human brain tissue with TPH1 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of wiDr cells using TPH1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Confocal immunofluorescent analysis of TPH1 Antibody (N-term)(Cat#AP6930a) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.