

CXCR-3 Polyclonal Antibody

Catalog # AP69344

Product Information

Application WB
Primary Accession P49682
Reactivity Human
Host Rabbit
Clonality Polyclonal
Calculated MW 40660

Additional Information

Gene ID 2833

Other Names CXCR3; GPR9; C-X-C chemokine receptor type 3; CXC-R3; CXCR-3; CKR-L2; G

protein-coupled receptor 9; Interferon-inducible protein 10 receptor; IP-10

receptor; CD antigen CD183

Dilution WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other

applications.

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CXCR3

Synonyms GPR9

Function [Isoform 1]: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11

and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G- protein signaling pathway (PubMed:12782716). Binds to CCL21. Probably promotes cell chemotaxis response. Upon activation by PF4, induces activated T-lymphocytes migration mediated via downstream Ras/extracellular signal-regulated kinase (ERK)

signaling. [Isoform 3]: Mediates the activity of CXCL11.

Cellular Location [Isoform 1]: Cell membrane; Multi-pass membrane protein


Tissue Location Isoform 1 and isoform 2 are mainly expressed in heart, kidney, liver and

skeletal muscle. Isoform 1 is also expressed in placenta. Isoform 2 is expressed in endothelial cells. Expressed in T-cells (at protein level).

Background

Isoform 1: Receptor for the C-X-C chemokine CXCL9, CXCL10 and CXCL11 and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G-protein signaling pathway (PubMed:12782716). Binds to CCL21. Probably promotes cell chemotaxis response. Isoform 3: Mediates the activity of CXCL11.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.