

# CYP2E1 Polyclonal Antibody

Catalog # AP69403

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">P05181</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	56849

## Additional Information

---

<b>Gene ID</b>	1571
<b>Other Names</b>	CYP2E1; CYP2E; Cytochrome P450 2E1; 4-nitrophenol 2-hydroxylase; CYPIIE1; Cytochrome P450-J
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---


<b>Name</b>	CYP2E1 {ECO:0000303   PubMed:10553002, ECO:0000312   HGNC:HGNC:2631}
<b>Function</b>	A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed: <a href="#">10553002</a> , PubMed: <a href="#">18577768</a> ). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed: <a href="#">10553002</a> , PubMed: <a href="#">18577768</a> ). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed: <a href="#">10553002</a> , PubMed: <a href="#">18577768</a> ). May be involved in the oxidative metabolism of xenobiotics (Probable).
<b>Cellular Location</b>	Endoplasmic reticulum membrane {ECO:0000250   UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250   UniProtKB:P05182}. Microsome membrane {ECO:0000250   UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250   UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250   UniProtKB:P05182}; Peripheral membrane protein

{ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}

## Background

Metabolizes several precarcinogens, drugs, and solvents to reactive metabolites. Inactivates a number of drugs and xenobiotics and also bioactivates many xenobiotic substrates to their hepatotoxic or carcinogenic forms.

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.