

CYP2S1 Polyclonal Antibody

Catalog # AP69406

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q96SQ9
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55817

Additional Information

Gene ID	29785
Other Names	CYP2S1; Cytochrome P450 2S1; CYP11B1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

Protein Information

Name	CYP2S1 {ECO:0000303 PubMed:11181079, ECO:0000312 HGNC:HGNC:15654}
Function	<p>A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:12711469, PubMed:21068195). In epidermis, may contribute to the oxidative metabolism of all-trans- retinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:12711469). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates (HPETEs) (PubMed:21068195). Independently of cytochrome P450 reductase, NADPH, and O2, catalyzes the breakdown of PGH2 to hydroxyheptadecatrienoic acid (HHT) and malondialdehyde (MDA), which is known to act as a mediator of DNA damage (PubMed:21068195).</p>
Cellular Location	Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein


Tissue Location

Expressed at higher levels in extrahepatic tissues including trachea, lung, stomach, small intestine, colon, kidney, breast, placenta and spleen (PubMed:11181079, PubMed:12711469) Expressed in peripheral blood leukocytes (PubMed:11181079) Constitutively expressed in skin (at protein level) (PubMed:12711469)

Background

Has a potential importance for extrahepatic xenobiotic metabolism.

Images



Western Blot analysis of various cells using CYP2S1 Polyclonal Antibody diluted at 1 : 2000

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.