

CysLTR2 Polyclonal Antibody

Catalog # AP69429

Product Information

ApplicationWB, IFPrimary AccessionQ9NS75ReactivityHumanHostRabbitClonalityPolyclonalCalculated MW39635

Additional Information

Gene ID 57105

Other Names CYSLTR2; CYSLT2; CYSLT2R; PSEC0146; Cysteinyl leukotriene receptor 2;

CysLTR2; G-protein coupled receptor GPCR21; hGPCR21; G-protein coupled

receptor HG57; HPN321

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000.

ELISA: 1/40000. Not yet tested in other applications. IF~~1:50~200

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium

azide.

Storage Conditions -20°C

Protein Information

Name CYSLTR2

Synonyms CYSLT2, CYSLT2R

Function Receptor for cysteinyl leukotrienes. The response is mediated via a G-protein

that activates a phosphatidylinositol-calcium second messenger system. Stimulation by BAY u9773, a partial agonist, induces specific contractions of pulmonary veins and might also have an indirect role in the relaxation of the

pulmonary vascular endothelium. The rank order of affinities for the

leukotrienes is LTC4 = LTD4 >> LTE4.

Cellular Location Cell membrane; Multi-pass membrane protein.

Tissue Location Widely expressed, with highest levels in the heart, placenta, spleen, peripheral


blood leukocytes and adrenal gland. In lung, expressed in the interstitial

macrophages, and slightly in smooth muscle cells

Background

Receptor for cysteinyl leukotrienes. The response is mediated via a G-protein that activates a phosphatidylinositol- calcium second messenger system. Stimulation by BAY u9773, a partial agonist, induces specific contractions of pulmonary veins and might also have an indirect role in the relaxation of the pulmonary vascular endothelium. The rank order of affinities for the leukotrienes is LTC4 = LTD4 >> LTE4.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.