

# Cystatin A Polyclonal Antibody

Catalog # AP69432

## Product Information

---

<b>Application</b>	WB, IHC-P, IF, ICC, E
<b>Primary Accession</b>	<a href="#">P01040</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	11006

## Additional Information

---

<b>Gene ID</b>	1475
<b>Other Names</b>	CSTA; STF1; STFA; Cystatin-A; Cystatin-AS; Stefin-A
<b>Dilution</b>	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200 ICC~~N/A E~~N/A
<b>Format</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
<b>Storage Conditions</b>	-20°C

## Protein Information

---

<b>Name</b>	CSTA
<b>Synonyms</b>	STF1, STFA
<b>Function</b>	This is an intracellular thiol proteinase inhibitor. Has an important role in desmosome-mediated cell-cell adhesion in the lower levels of the epidermis.
<b>Cellular Location</b>	Cytoplasm.
<b>Tissue Location</b>	Expressed in the skin throughout the epidermis.

## Background

---

This is an intracellular thiol proteinase inhibitor. Has an important role in desmosome-mediated cell-cell adhesion in the lower levels of the epidermis.

## Images

---



Western Blot analysis of various cells using Cystatin A Polyclonal Antibody



Western Blot analysis of HT29 cells using Cystatin A Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.