

TH Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP6945b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	P07101
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58600
Antigen Region	486-514

Additional Information

Gene ID	7054
Other Names	Tyrosine 3-monooxygenase, Tyrosine 3-hydroxylase, TH, TH, TYH
Target/Specificity	This TH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 486-514 amino acids from the C-terminal region of human TH.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TH Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TH (HGNC:11782)
Synonyms	TYH
Function	Catalyzes the conversion of L-tyrosine to L- dihydroxyphenylalanine (L-Dopa), the rate-limiting step in the biosynthesis of catecholamines, dopamine, noradrenaline, and adrenaline. Uses tetrahydrobiopterin and

molecular oxygen to convert tyrosine to L-Dopa (PubMed:[15287903](#), PubMed:[1680128](#), PubMed:[17391063](#), PubMed:[24753243](#), PubMed:[34922205](#), PubMed:[8528210](#), Ref.18). In addition to tyrosine, is able to catalyze the hydroxylation of phenylalanine and tryptophan with lower specificity (By similarity). Positively regulates the regression of retinal hyaloid vessels during postnatal development (By similarity).

Cellular Location

Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P24529}. Nucleus {ECO:0000250|UniProtKB:P04177} Cell projection, axon {ECO:0000250|UniProtKB:P24529}. Cytoplasm {ECO:0000250|UniProtKB:P04177}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:P04177}. Note=When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals. {ECO:0000250|UniProtKB:P04177}

Tissue Location

Mainly expressed in the brain and adrenal glands.


Background

TH is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons.


References

Kuhn,D.M., et.al., J. Biol. Chem. 277 (16), 14336-14342 (2002)


Images



TH Antibody (C-term) (Cat. #AP6945b) western blot analysis in SH-SY5Y cell line lysates (35ug/lane). This demonstrates the TH antibody detected the TH protein (arrow).



Formalin-fixed and paraffin-embedded human brain tissue with TH Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of K562 cells using TH Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.