

# DACA-1 Polyclonal Antibody

Catalog # AP69466

## Product Information

Application	WB, IHC-P
Primary Accession	<a href="#">Q9BYJ9</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	60874

## Additional Information

Gene ID	54915
Other Names	YTHDF1; C20orf21; YTH domain family protein 1; Dermatomyositis associated with cancer putative autoantigen 1; DACA-1
Dilution	WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.
Storage Conditions	-20°C

## Protein Information

Name	YTHDF1 {ECO:0000303 Ref.4, ECO:0000312 HGNC:HGNC:15867}
Function	Specifically recognizes and binds N6-methyladenosine (m6A)- containing mRNAs, and regulates their stability (PubMed: <a href="#">24284625</a> , PubMed: <a href="#">26318451</a> , PubMed: <a href="#">32492408</a> , PubMed: <a href="#">39900921</a> ). M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in mRNA stability and processing (PubMed: <a href="#">24284625</a> , PubMed: <a href="#">32492408</a> ). Acts as a regulator of mRNA stability by promoting degradation of m6A-containing mRNAs via interaction with the CCR4-NOT complex (PubMed: <a href="#">32492408</a> ). The YTHDF paralogs (YTHDF1, YTHDF2 and YTHDF3) shares m6A-containing mRNAs targets and act redundantly to mediate mRNA degradation and cellular differentiation (PubMed: <a href="#">28106072</a> , PubMed: <a href="#">32492408</a> ). Required to facilitate learning and memory formation in the hippocampus by binding to m6A-containing neuronal mRNAs (By similarity). Acts as a regulator of axon guidance by binding to m6A-containing ROBO3 transcripts (By similarity). Acts as a negative regulator of antigen cross-presentation in myeloid dendritic cells (By similarity). In the context of tumorigenesis, negative regulation of antigen cross-presentation limits the anti-tumor response by reducing efficiency of tumor-antigen cross- presentation (By similarity). Promotes

formation of phase-separated membraneless compartments, such as P-bodies or stress granules, by undergoing liquid-liquid phase separation upon binding to mRNAs containing multiple m6A-modified residues: polymethylated mRNAs act as a multivalent scaffold for the binding of YTHDF proteins, juxtaposing their disordered regions and thereby leading to phase separation (PubMed:[31292544](#), PubMed:[31388144](#), PubMed:[32451507](#)). The resulting mRNA-YTHDF complexes then partition into different endogenous phase-separated membraneless compartments, such as P-bodies, stress granules or neuronal RNA granules (PubMed:[31292544](#)).


#### Cellular Location

Cytoplasm. Cytoplasm, P-body. Cytoplasm, Stress granule

## Background

Specifically recognizes and binds N6-methyladenosine (m6A)-containing mRNAs, and promotes mRNA translation efficiency (PubMed:[24284625](#), PubMed:[26046440](#), PubMed:[26318451](#)). M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in the efficiency of mRNA splicing, processing and stability (PubMed:[24284625](#)). Acts as a regulator of mRNA translation efficiency: promotes ribosome loading to m6A-containing mRNAs and interacts with translation initiation factors eIF3 (EIF3A or EIF3B) to facilitate translation initiation (PubMed:[26046440](#)).

## Images



Western Blot analysis of various cells using DACA-1 Polyclonal Antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.